

RCF 1

Theories of PR Maps and Partial PR Maps

$\widehat{\mathbf{PR}}_{\mathbf{A}}$

Michael Pfender*

May 2008[†]

Abstract: We give to the categorical theory $\mathbf{PR}$ of Primitive Recursion a logically simple, *algebraic* presentation, via *equations between maps*, plus one genuine HORNER type schema, namely Freyd’s uniqueness of the *initialised iterated*. Free Variables are introduced – formally – as another names for *projections*. Predicates $\chi : A \rightarrow 2$ admit *interpretation* as (formal) Objects $\{A | \chi\}$ of a surrounding Theory $\mathbf{PR}_{\mathbf{A}} = \mathbf{PR} + (\text{abstr}) : \text{schema} (\text{abstr})$ formalises this *predicate abstraction* into additional Objects. Categorical Theory $\mathbf{PR}_{\mathbf{A}} \sqsupset \mathbf{PR}_{\mathbf{A}} \sqsupset \mathbf{PR}$ then is the Theory of formally *partial* PR-maps, having Theory $\mathbf{PR}_{\mathbf{A}}$ embedded. This Theory $\widehat{\mathbf{PR}}_{\mathbf{A}}$ bears the structure of a (still) diagonal monoidal category. It is equivalent to “the” categorical theory of μ -recursion (and of while loops), viewed as partial PR maps. So the present approach to partial maps sheds new light on CHURCH’s THESIS, “embedded” into a Free-Variables, formally *variable-free* (categorical) framework.

⁰ This is part 1 of a cycle on *Recursive Categorical Foundations*.
There is a still more detailed version (pdf file) equally entitled *Theories of PR Maps and Partial PR Maps*

⁰ Legend of LOGO: $\widehat{\mathbf{PR}}_{\mathbf{A}} = \widehat{\mathbf{PR}}_{\mathbf{A}}$ symbolises the theory of *partial* maps over categorical Free-Variables theory $\mathbf{PR}_{\mathbf{A}}$ of Primitive Recursion, $\mathbf{PR}$, with predicate Abstraction

*TU Berlin, Mathematik, pfender@math.tu-berlin.de

[†]last revised September 22, 2008

1 Introduction

We develop here, from scratch, a formally *variable-free*, *categorical* Theory of μ -*recursion*, without use of formal *quantification*: This Theory is formalised on the basis of a theory $\widehat{\mathbf{PR}}_{\mathbf{A}}$ of *partial* PR maps which in turn is introduced as a *definitional*, conservative extension of Theory $\mathbf{PR}_{\mathbf{A}} = \mathbf{PR} + (\text{abstr})$, the latter obtained from *fundamental* categorical Theory $\mathbf{PR}$ of Primitive Recursion: we formally interpret PR *predicates* $\chi : A \rightarrow 2$ as additional, *defined* Objects of emerging Theory $\mathbf{PR}_{\mathbf{A}}$: schema of abstraction already “hidden” in *fundamental* Theory $\mathbf{PR}$ of Primitive Recursion. The latter is given as *Cartesian Hull* over data and axioms of a Natural Numbers Object $\mathbb{N}$, this in the sense of LAWVERE, EILENBERG & ELGOT, and FREYD.

Central in present approach is the notion of a *partial* PR *map*: Such a (formally) partial map $f : A \multimap B$ is given, in categorical, variable-free terms, by:

- an *enumeration source* Object D_f in $\mathbf{PR}_{\mathbf{A}}$ (e.g.: $D_f = \mathbb{N}$),
- a $\mathbf{PR}_{\mathbf{A}}$ -map $d_f : D_f \rightarrow A$, meaning for $\mathbf{PR}_{\mathbf{A}}$ -*enumeration* of *defined arguments* of f ,
- and a $\mathbf{PR}_{\mathbf{A}}$ -*rule*-map $\widehat{f} : D_f \rightarrow B$ for f ,
- these data with (intuitive) meaning: for $a \in A$ *defined argument*, of form $a = d_f(\hat{a}) \in A$ for suitable $\hat{a} \in D_f$: $d_f(\hat{a}) = a \xrightarrow{f} \widehat{f}(\hat{a}) \in B$.

We prove a **Structure Theorem** for *partial-map extensions* $\widehat{\mathbf{S}} \sqsupset \mathbf{S}$, where $\mathbf{S}$ is a Cartesian PR theory with schema of predicate abstraction – mostly $\mathbf{S} := \mathbf{PR}_{\mathbf{A}}$ or one of its strengthenings – which establishes these extensions (via embedding) as *diagonal monoidal* PR theories: Cartesian structure is lost in part, since the (still present) projection- and terminal-map-families do not preserve their character as *natural transformations* in the extension. These *partial-map extensions* turn out to be *Closures*: $\widehat{\widehat{\mathbf{S}}} \cong \widehat{\mathbf{S}} : \text{Partial}$ partial maps have a representation as just *partial maps*.

Within this Free-Variables (formally: variable free) categorical framework $\widehat{\mathbf{PR}}_{\mathbf{A}}$ for *partial* PR map theories, we discuss (Free-Variables) *category based* μ -recursion as well as *content driven* loops such as while loops.

This prepares in particular discussion of *termination* for suitable special such loops, namely those given as *Complexity Controlled Iterations*, for which iteration *step* decreases a *complexity* measure within a suitably given (constructive) *ordinal* O , “until” minimum 0 of O is reached. Complexity Controlled Iteration is basic for the following second part of investigation on *Recursive Categorical Foundations*, RCF 2, entitled *Evaluation and Consistency*.

Evaluation of PR map codes is there *resolved* into such an iteration with *descending* complexity values. “Hence” we can “hope” this formally partial evaluation to always *terminate*, by reaching complexity 0. In that case within ordinal O taken the lexicographically ordered set of polynomials over $\mathbb{N}$, in one indeterminate.

2 Notions, Axioms, Results for Theories **PR** and **PR_A**

Fundamental Theory **PR** of *Primitive Recursion* is the *minimal*, “*initial*” *Cartesian* Theory with (universal) Natural Numbers Object:


As Objects it has $\mathbb{1}, N, \dots, A, B, \dots, (A \times B)$, i. e. all (binary bracketed) finite powers of Natural Numbers Object (“NNO”) $\mathbb{N}$.

It comes with associative *composition* “ $\circ$ ”, functorial *cylindrification* $(A \times g) : A \times B \rightarrow A \times B'$ – and from this with bifunctorial *Cartesian product* “ $\times$ ” – and is generated over *basic map constants* $0 : \mathbb{1} \rightarrow \mathbb{N}$ (*zero*), *successor* $s : \mathbb{N} \times \mathbb{N}$, as well as *terminal map* $! : A \rightarrow \mathbb{1}$, *diagonal* $\Delta : A \rightarrow A \times A$, and *projections* $\ell : A \times B \rightarrow A$ and $r : A \times B \rightarrow B$.

For given $f : C \rightarrow A$ and $g : C \rightarrow B$, *induced map* $(f, g) : C \rightarrow A \times B$ into Cartesian Product is **defined** as

$$(f, g) =_{\text{def}} (f \times g) \circ \Delta_C : C \rightarrow C^2 =_{\text{by def}} C \times C \rightarrow A \times B,$$


with $(f \times g) : C^2 \rightarrow A \times B$ defined by either – equal(!) – sequence of cylindrified above, see diamond sub-DIAGRAM in



Binary Cartesian Map Product DIAGRAM

In present context we take GODEMENT’s equations for Cartesian (!) Prod-


uct, as **axioms**:



Uniqueness of the induced $(f, g) \stackrel{\text{by def}}{=} (f \times g) \circ \Delta_C : C \rightarrow C^2 \rightarrow A \times B$ is *forced* – **axiomatically** – by the following FOURMAN’s equation (equational *schema*):

$$\text{(Four!)} \quad \frac{h : C \rightarrow A \times B \text{ map}}{(\ell_{A,B} \circ h, r_{A,B} \circ h) = h : C \rightarrow A \times B.}$$

The NNO $\mathbb{1} \xrightarrow{0} \mathbb{N} \xrightarrow{s} \mathbb{N}$ is given – by **axiom** – the following property, expressed as *iteration schema*:




Basic Iteration DIAGRAM

As Uniqueness schema we need the following one – of FREYD, formally stronger than *uniqueness of the iterated* $f^s : A \times \mathbb{N} \rightarrow A$ satisfying the equations of Basic Iteration DIAGRAM above – namely **Uniqueness of initialised iterated**:

$$\begin{array}{l} f : A \rightarrow B, g : B \rightarrow B, h : A \times \mathbb{N} \rightarrow B, \text{ all in } \mathbf{PR}, \\ h \circ (\text{id}_A, 0 \circ !_A) = f : A \rightarrow B, \text{ (init)} \\ h \circ (A \times s) = g \circ h : A \times \mathbb{N} \rightarrow B, \text{ (step)} \\ \hline \text{(FR!)} \quad h = g^s \circ (f \times \mathbb{N}) : A \times \mathbb{N} \rightarrow B \times \mathbb{N} \rightarrow B, \end{array}$$

in terms of FREYD’s pentagonal diagram:



FREYD's uniqueness DIAGRAM (FR!)

Schemata (it) and (FR!) give in fact the well-known full schema (pr) of Primitive Recursion:

$$\begin{array}{l}
 \text{(pr)} \quad \frac{
 \begin{array}{l}
 g = g(a) : A \rightarrow B \text{ PR } (init \text{ map}) \\
 h = h((a, n), b) : (A \times \mathbb{N}) \times B \rightarrow B \text{ (step map)}
 \end{array}
 }{
 \begin{array}{l}
 f = f(a, n) = \text{pr}[g, h](a, n) : A \times \mathbb{N} \rightarrow B \text{ in } \mathbf{PR} \text{ such that} \\
 f(a, 0) = g(a) : A \rightarrow B \text{ (init), and} \\
 f(a, s \ n) = h((a, n), f(a, n)) : (A \times \mathbb{N}) \rightarrow B, \text{ (step)} \\
 \text{as well as} \\
 \text{(pr!) : } f \text{ is unique with these properties.}
 \end{array}
 }
 \end{array}$$

This *full schema of PR* has as consequence in particular the elegant and powerfull **Uniqueness schemata** of GOODSTEIN, $U_1 - U_4$, which *passive* parameter, usually $a \in \mathbb{N}$ here made explicit.

Using these schemata, GOODSTEIN proves central equations for *addition*, *truncated subtraction*, and *multiplication* on the “NNO” $\mathbb{N}$:

As usual, the basic structure of a *unitary commutative semiring* on NNO $\mathbb{N}$ – (plus *truncated subtraction and exponentiation*) – is **defined** – and **characterised**, map theoretically – with interpretation of Free Variables a and n as

projections $a := \ell : A \times \mathbb{N} \rightarrow A$ and $n := r : A \times \mathbb{N} \rightarrow \mathbb{N}$ – as follows:

$0, 1 \stackrel{\text{by def}}{=} s\,0 : \mathbb{1} \rightarrow \mathbb{N}$: *zero* resp. *one*,
addition $a + n = s^{\S}(a, n) : \mathbb{N}^2 \rightarrow \mathbb{N}$ by
 $a + 0 = a : \mathbb{N} \rightarrow \mathbb{N}$, $a + s\,n = s(a + n) : \mathbb{N}^2 \rightarrow \mathbb{N}$,
truncated subtraction $a \dot{-} n : \mathbb{N}^2 \rightarrow \mathbb{N}$ by:
 $a \dot{-} 0 = a$, $a \dot{-} s\,n = \text{pre}(a \dot{-} n)$,
 with *predecessor* $\text{pre} : \mathbb{N} \rightarrow \mathbb{N}$
 PR defined (and characterised) by $\text{pre}\,0 = 0$, $\text{pre}\,s\,n = n$,
 so for example $5 \dot{-} 2 = 3$, but $2 \dot{-} 5 = 0$;
multiplication $a \cdot n : \mathbb{N}^2 \rightarrow \mathbb{N}$ by
 $a \cdot 0 = 0$, $a \cdot s\,n = a \cdot n + a$,
 and *exponentiation* $a^n : \mathbb{N}^2 \rightarrow \mathbb{N}$ by
 $a^0 = 0$, $a^{s\,n} = a^n \cdot a$.

Remark on use of free variables: Interpretation of Free Variables as (identities) resp. *projections*, (possibly nested) is subject to formal rules extending this interpretation in the above example. Vice versa, projections can be seen as free variables, by (re)naming them with names $a, b, \dots, x, y, \dots$ usually standing for free “individual” variables. From now on, we will make extensive use of Free Variables, *anchored* in the Cartesian structure of *basic* theory $\mathbf{PR}_A = \mathbf{PR} + (\text{abstr})$ and its strengthenings.

Continuation of *elementary map equations* for NNO $\mathbb{N} : \text{Multiplication}$ on $\mathbb{N}$ *distributes* not only over addition but over *truncated subtraction* as well, almost (again) by definition.

(Boolean) *Logic* and *Order* then are defined, and characterised as follows, the latter essentially via truncated subtraction:

$\text{sign} = \text{sign}(n) : \mathbb{N} \rightarrow \mathbb{N}$ by $\text{sign}(0) = 0$, $\text{sign}(s\,n) = 1 = s\,0 : \mathbb{N} \rightarrow 2$,
 $\neg = \neg(n) : \mathbb{N} \rightarrow \mathbb{N}$ by $\neg(0) = 1$, $\neg(s\,n) = 0$.

A map $\chi : A \rightarrow \mathbb{N}$, here such a map within $\mathbf{PR}$, is called a *predicate* (on Object A) if

$$\text{sign} \circ \chi = \neg \circ \neg \circ \chi = \text{chi} : \mathbb{N} \rightarrow \mathbb{N}.$$

By obvious reason, we write for this

$$\chi : A \rightarrow 2 \subset \mathbb{N}.$$

For the moment, this is just notation. Later, in theory $\mathbf{PR}_A$ to come, Object $2 = \{n \in \mathbb{N} \mid n < s\,0\} \subset \mathbb{N}$ will become an *Object* on its own right.¹

¹ The idea to introduce Boolean (!) Free-Variables predicate Calculus into the Theory of Primitive Recursion this way, without an explicit basic (“undefined”) Object $2 \equiv \{\text{false}, \text{true}\} \equiv \{0, 1\}$ goes back to GOODSTEIN 1971 and REITER 1982.

Using this notation, we get the Boolean Operations, and basic binary predicates on $\mathbb{N}$ as

$$\begin{aligned} \wedge &= [a \wedge b] =_{\text{def}} \text{sign}(a \cdot b) : \mathbb{N}^2 \rightarrow \mathbb{N}, \\ [m \leq n] &=_{\text{def}} \neg [m \dot{-} n] : \mathbb{N}^2 \rightarrow 2, \\ [m < n] &=_{\text{def}} \text{sign}[n \dot{-} m] : \mathbb{N}^2 \rightarrow 2, \\ &\text{and directly from “}\leq\text{”}: \\ [m \dot{=} n] &= [m \leq n] \wedge [n \leq m] : \mathbb{N}^2 \rightarrow 2. \end{aligned}$$

The latter predicate is *predicative equality*. It can be extended to all *fundamental Objects* (finite bracketed powers of $\mathbb{N}$), by componentwise conjunctive definition. And furthermore to (formal, virtual) predicate *extensions* $\{A \mid \chi : A \rightarrow 2\}$ of **PR**-predicates χ , see below.

[Addition as well as multiplication – the latter with arguments greater zero – are strictly *monotonous* with respect to the (linear) order introduced above via truncated subtraction.]

This *equality “of individuals”* $[a \dot{=} a'] : A^2 \rightarrow 2$ is *reflexive, symmetric, and transitive*. On $\mathbb{N}$ it satisfies – with regard to strict order $< : \mathbb{N}^2 \rightarrow \mathbb{N}$ – the *law of trichotomy*.

Furthermore it is – as we expect it from a “sound” notion of *equality* – *substitutive* (in LEIBNIZ’ sense), with respect to the “basic meta-operations” *composition, Cartesian Product*, as well as *Iteration*, and therefore in particular with respect to *all* operations (above), introduced on $\mathbb{N}$ via these basic meta-operations.

The key relationship between this *predicate* and the – already given – notion of equality between maps:

$$\begin{aligned} f &=^{\mathbf{PR}} g : A \rightarrow B, \text{ also written} \\ \mathbf{PR} &\vdash f = g : A \rightarrow B : \\ \text{Theory } \mathbf{PR} &\text{ derives } \underline{\text{equality}} \ f = g, \end{aligned}$$

is the following *Equality Definability Schema*, which is derivable in theory **PR**, and also in Theory **PR_A** = **PR**+(abstr) to come, as well as in all strengthenings of these:

Equality Definability Theorem: An Arithmetical Theory **T**, i. e. **T** an extension of **PR**, admits the following schema (EquDef):

$$\begin{aligned} &f, g : A \rightarrow \mathbb{N} \text{ in } \mathbf{T}, \\ (\text{EquDef}) \quad &\frac{\mathbf{T} \vdash \dot{\circ}(f, g) = \text{true}_{\mathbb{N}^2} : \mathbb{N}^2 \rightarrow 2}{\mathbf{T} \vdash f = g : A \rightarrow \mathbb{N}, \text{ algebraically:}} \\ &f =^{\mathbf{T}} g : A \rightarrow \mathbb{N}. \end{aligned}$$

Equality Definability extends to **T**-map pairs $f, g : A \rightarrow B$ with (common) codomain a (finite) Cartesian Product B of Objects $\mathbb{N}$, a *fundamental* Object

– in **PR** – or even B a predicative extension $B = \{C \mid \gamma\}$ of such an Object, an Object of Theory **PR**_A for short, see below.

Remark: For **proof** of the laws of multiplication, and also for proof of *logically* all important *Equality Definability* above, GOODSTEIN proves commutativity of the maximum function, namely

$$\max(a, b) \stackrel{\text{def}}{=} a + (b \dot{-} a) = b + (a \dot{-} b) \stackrel{\text{by def}}{=} \max(b, a) : \mathbb{N}^2 \rightarrow \mathbb{N}.$$

We now “realise” straight forward the schema of *predicate abstraction*: We start with a (**PR**) predicate $\chi : A \rightarrow 2$ on a *fundamental* Object $A : \text{Object}$ A is a (binary bracketed) Cartesian product out of $\mathbb{N}$ and $\mathbb{1}$. Such a predicate $\chi : A \rightarrow 2$, formally: $\chi : \mathbb{N} \rightarrow \mathbb{N}$ (see above), can be turned into a *virtual*, “new” Object $\{A \mid \chi\} = \{a \in A \mid \chi(a)\}$ easily: just take as Objects of *new frame*, extended Theory **PR**_A the fundamental maps of form $\chi : A \rightarrow 2$ (*predicates*) within *fundamental* Theory **PR** above, and as maps between such Objects $\{A \mid \chi\}$ and $\{B \mid \psi\}$ those **PR**-maps $f : A \rightarrow B$ which transform χ into ψ . Two such maps $f, g : A \rightarrow B$ are *identified*, declared *equal*, if they agree “on” χ . This *definitional, conservative* extension is called here **PR**_A = **PR** + (abstr) $\sqsupset$ **PR**, the *basic Theory of Primitive Recursion*.

Structure Theorem for PR_A : *Basic* Theory **PR**_A becomes a *Cartesian Category* (Theory), has all *Extensions* of its predicates and therefore all *equalisers*, and contains *fundamental* Theory **PR** embedded via

$$\langle f : A \rightarrow B \rangle \mapsto \langle f : \{A \mid \text{true}_A\} \rightarrow \{B \mid \text{true}_B\} \rangle.$$

Furthermore, it has all equality predicates – by restriction – and admits the schema of *Equality Definability*.

Proof (REITER 1980): A preliminary version of **PR**_A is constructed as canonical extension of **PR**’s Class of Objects into its predicates and then admitting as *maps* between these “new” Objects those of **PR** which are compatible with the given, *defining* predicates for these “Objects”. This way one gets a Cartesian PR theory. Equality of **PR**_A maps, given by their equality on the predicate of their **PR**_A Domain Object, is compatible with this Cartesian PR structure and thereby gives the Cartesian PR structure of **PR**_A, defined as Quotient theory by this *notion of equality*. Forming the *iterated* is likewise compatible with this canonical notion of equality in the extended world, and therefore **PR**_A becomes this way a PR category. By construction, **PR**_A gets in addition the wanted predicate-extension and equaliser-by-extension structure: it has finitie limits, in particular pullbacks (and multiple pullbacks) **q.e.d.**

Remarks:

(i) A **PR**_A-map $f : \{A \mid \chi\} \rightarrow \{B \mid \psi\}$ can be viewed as a *defined partial PR* map from A to B with values in $\psi : \text{Set of defined arguments}$, namely $\{a \in A \mid \chi(a)\}$ is PR *decidable*. By **definitin** of **PR**_A’s equality, **PR**-map $f : A \rightarrow B$ “doesn’t care” about arguments a in the *complement* $\{a \in A \mid \neg \chi(a)\}$.

So wouldn't it be easier to realise this view to *defined partial maps* just by throwing the *undefined arguments* into a *waste basket* $\{\perp\}$ say?

But where to place this waste basket, this for each Codomain Object B ? The fundamental Objects have a zero-vector as a candidate. For example we could interpret truncated subtraction as a *defined partial map*

$$a \dot{-} b : \{(m, n) \in \mathbb{N} \times \mathbb{N} \mid m \geq n\} \rightarrow \mathbb{N},$$

and throw the complement $\{(m, n) \in \mathbb{N} \times \mathbb{N} \mid m < n\}$ into waste basket $\{0\} \subset \mathbb{N}$. But this is not a good interpretation of *truncated* (!) subtraction: Value 0 is *not* waste, it has an important meaning as zero.

“The” waste basket $\{\perp\}$ should be an entity with a *natural* extra representation, and we should have only one such entity in a later theory of defined partial PR maps to come. This theory, to be called $\mathbf{PR}_A^{\mathbb{X}}$, will be constructed with the help of *Universal Object* $\mathbb{X}$ which is to contain *codes* of all singletons and (nested) pairs of natural numbers, and “below” these codes it has room for code of *undefined value* symbol $\perp$, in a “Hilbert’s hotel”. All this will be carried through within present theory $\mathbf{PR}_A$.

(ii) A $\mathbf{PR}$ -map $f : A \rightarrow B$ such that f “is” a $\mathbf{PR}_A$ -map

$$\begin{aligned} f : \{A \mid \chi \vee \chi' : A \rightarrow 2\} &\rightarrow \{B \mid \psi\}, \text{ also “works” as a } \mathbf{PR}_A\text{-map} \\ f : \{A \mid \chi\} &\rightarrow \{B \mid \psi\}, \text{ and a } \mathbf{PR}_A\text{-map} \\ g : \{A \mid \chi\} &\rightarrow \{B \mid \psi \wedge \psi'\} \text{ also “works” as a } \mathbf{PR}_A\text{-map} \\ g : \{A \mid \chi\} &\rightarrow \{B \mid \psi\}. \end{aligned}$$

Since map-properties of *injectivity*, *epi-property* of $\mathbf{PR}$ -maps viewed as $\mathbf{PR}_A$ -maps, **depend** on choice of hosting (predicative) $\mathbf{PR}_A$ Objects – **examples** above – *specification* of a $\mathbf{PR}_A$ map $f : \{A \mid \chi\} \rightarrow \{B \mid \psi\}$ must contain, besides $\mathbf{PR}$ -map $f : A \rightarrow B$, Domain and Codomain *Objects* $\chi : A \rightarrow 2$ and $\psi : B \rightarrow 2$ as well. This way the members of Map set family $\mathbf{PR}_A(A, B) : A, B \mathbf{PR}_A$ -Objects, become mutually disjoint. Inclusions $i : A' \xrightarrow{\subseteq} A''$ are realised in $\mathbf{PR}_A$ as restricted $\mathbf{PR}$ -identities $\text{id}_A : \{A \mid \chi'\} \xrightarrow{\subseteq} \{A \mid \chi''\}$, $\chi' \implies \chi''$.

3 Goodstein Arithmetic **GA**

In “Development of Mathematical Logic” (Logos Press 1971) R. L. Goodstein gives four basic uniqueness-rules for Free-Variable Arithmetics. We show here that these four rules are sufficient for proving the commutative and associative laws for multiplication and the distributive law, for addition as well as for truncated subtraction.

We include² into Goodstein’s uniqueness-rules a “passive parameter” a . These extended rules are provable using Freyd’s uniqueness Theorem (pr!), part

²Sandra Andrasko and the author

of *full schema* (pr) of Primitive Recursion which he deduces from his uniqueness (FR!) of the *initialised iterated*. FREYD deduces the latter from availability of a Natrual Numbers Object $\mathbb{N}$ in LAWVERE's sense, and (!) *axiomatic* availability of “higher order” *internal* hom objects with, again axiomatic, *evaluation* map family for these objects, of form $\epsilon_{A,B} : B^A \times A \rightarrow B$ within (!) the category considered.

Goodstein's rules with passive parameter: Let $f, g : A \times N \rightarrow N$ be primitive recursive maps, $s : N \rightarrow N$ the successor map $n \mapsto n + 1$ and $\text{pre} : N \rightarrow N$ the predecessor map, usually written as $n \mapsto n \dot{-} 1$.

Then Goodstein's rules read:

$$\begin{array}{l}
\text{U}_1 \quad \frac{f(a, sn) = f(a, n) : A \times \mathbb{N} \rightarrow B}{f(a, n) = f(a, 0) : A \times \mathbb{N} \rightarrow B} \\
\text{U}_2 \quad \frac{f(a, sn) = s f(a, n) : A \times \mathbb{N} \rightarrow \mathbb{N}}{f(a, n) = f(a, 0) + n : A \times \mathbb{N} \rightarrow \mathbb{N}} \\
\text{U}_3 \quad \frac{f(a, sn) = \text{pre } f(a, n) : A \times \mathbb{N} \rightarrow \mathbb{N}}{f(a, n) = f(a, 0) \dot{-} n : A \times \mathbb{N} \rightarrow \mathbb{N}} \\
\text{U}_4 \quad \frac{\begin{array}{l} f(a, 0) = g(a, 0) : A \rightarrow \mathbb{N} \\ f(a, sn) = g(a, sn) : A \times \mathbb{N} \rightarrow \mathbb{N} \end{array}}{f(a, n) = g(a, n) : A \times \mathbb{N} \rightarrow \mathbb{N}}
\end{array}$$

Comment: Theories **PR** and **PR_A** allow, within rules U_1 and U_4 above, for replacing $\mathbb{N}$ as a Codomain Object, by an arbitrary object B of **PR** resp. **PR_A**.

Rule U_4 , of *uniqueness* of maps defined by **case distinction**, is nothing else than *uniqueness* of the *induced map out of the sum* $A \times \mathbb{N} \cong (A \times \mathbb{1}) + (A \times \mathbb{N})$, this sum canonically realised via *injections* $\iota = (\text{id}_A \times 0) : A \times \mathbb{1} \rightarrow A \times \mathbb{N}$ as well as, right injection: $\kappa = \text{id}_A \times s : A \times \mathbb{N} \rightarrow A \times \mathbb{N}$:

This uniqueness combined with LEIBNIZ' *compatibility of induced-map-out-of-a-sum* with map (term) equality, compatibility available in Theories **PR**, **PR_A**, and their strengthenings.

Proof of these four rules is straight forward for theories **PR**, **PR_A** (and strengthenings), using FREYD's uniqueness (FR!) and uniqueness clause (pr!) of the *full schema of Primitive Recursion* respectively.

U_1 - U_4 give, by means of a derived schema V_4 , the wanted

Structure Theorem for NNO $\mathbb{N} : \mathbb{N}$ admits the structure of a *commutative semiring* with zero, *truncated subtraction* $\dot{-} = m \dot{-} n \rightarrow \mathbb{N}$, over which multiplication *distributes*, a *linear order* $m < n : \mathbb{N}^2 \rightarrow 2$, and *equality predicate* $\dot{=} = [m \dot{-} n] : \mathbb{N}^2 \rightarrow 2$, both of the latter **defined** via truncated subtraction.

Order and equality (predicates) satisfy the *law of trichotomy*; addition is strictly monotoneous in both arguments; truncated subtraction is weakly monotoneous in first, and weakly antitoneous in second argument, whereas multiplication is strongly monotoneous in both arguments, on $\mathbb{N}_{>0}^2 = \{\mathbb{N} \mid > 0\}^2$.

$\mathbb{N}$ and $\{n \in \mathbb{N} \mid n < 2\}$ admit (2-valued) **Boolean Logic** sign, $\neg$, $\wedge$, $\vee$, $\implies$.

Last but not least, the maximum $\max(a, b) : \mathbb{N}^2 \rightarrow \mathbb{N}$ *commutes*:


$$\mathbf{PR} \vdash \max(a, b) \stackrel{\text{def}}{=} a + (b \dot{-} a) = b + (a \dot{-} b) \stackrel{\text{by def}}{=} \max(b, a) : \mathbb{N}^2 \rightarrow \mathbb{N}.$$

This goes into **Proof of Equality Definability**, for **PR**, **PR_A**, and strengthenings.

Proof, by $U_1 - U_4$, straightforward but tedious.

4 Theories of Partial PR Maps

We now turn to Extension of PR theories – here Theory **PR_A** or a strengthening – into Theories of *partial* PR maps (*map terms*). This extension is understood best, wenn looking at the following **DIAGRAM** which shows *composition* of two such partial maps, $g : B \rightarrow C$ with $f : A \rightarrow B$:



Composition **DIAGRAM** for $\widehat{\mathbf{S}}$

A partial map $f = \langle (d_f : \widehat{f}) : D_f \rightarrow A \times B \rangle : A \rightarrow B$ consists of a (PR) enumeration $d_f = d_f(\hat{a}) : D_f \rightarrow A$ of *defined-arguments* for f , and a *rule* $\widehat{f} = \widehat{f}(\hat{a}) : D_f \rightarrow B$, fixing the values $f(a) \stackrel{\text{def}}{=} \widehat{f}(\hat{a})$ for *defined* arguments $a \in A$, i. e. a of form $a = d_f(\hat{a})$, $\hat{a} \in D_f$.

Up to here, f defines just a *relation*, in the sense of BRINKMANN/PUPPE 1969. What is lacking is *right uniqueness*, see the following

Partial-Map Schema:


$$\begin{array}{l}
\gamma f = \gamma f(\hat{a}) : D_f \rightarrow A \times B \text{ } \mathbf{S}\text{-map,} \\
\text{called } \textit{graph} \text{ (of } f : A \rightarrow B \text{ to be introduced),} \\
d_f = d_f(\hat{a}) =_{\text{def}} \ell \circ \gamma f : D_f \rightarrow A \text{ } \textit{defined arguments enumeration} \\
\hat{f} = \hat{f}(\hat{a}) : D_f \rightarrow B \text{ } \textit{rule} \\
\mathbf{S} \vdash d_f(\hat{a}) \doteq d_f(\hat{a}') \implies \hat{f}(\hat{a}) \doteq \hat{f}(\hat{a}') : D_f^2 \rightarrow 2 \text{ } (\hat{a}, \hat{a}' \in D_f \text{ free}) : \\
\textit{right uniqueness} \\
(\widehat{\mathbf{S}}) \quad \hline \\
f =_{\text{def}} \langle \gamma f = (d_f, \hat{f}) : D_f \rightarrow A \times B \rangle : A \rightarrow B \\
\widehat{\mathbf{S}}\text{-morphism, } \textit{partial } \mathbf{S} \text{ map.}
\end{array}$$

DIAGRAM above then shows – BRINKMANN & PUPPE type – *partial map composition* $g \hat{\circ} f : A \rightarrow B \rightarrow C$ via pullback.

Equality $f \hat{=} g : A \rightarrow B$ of partial maps – over $\mathbf{S}$ – is established by availability of a pair $D_f \xrightleftharpoons[i]{i} D_{f'}$ of *defined-arguments comparison* maps (in $\mathbf{S}$)

which are compatible as such with $d_f, d_{f'}$ as well as with $\hat{f}, \hat{f}'$. Availability of just one of these $\mathbf{S}$ maps, of $i : D_f \rightarrow D_{f'}$ say, **defines** “graph” *inclusion*, here $f \hat{\subseteq} f' : A \rightarrow B$.

Basic *compatibility* of (partial) *composition* “ $\hat{\circ}$ ”, with graph *inclusion* $\hat{\subseteq}$ – and hence with *partial equality* “ $\hat{=}$ ” then is given by the universal properties of (composed) pullback in the following DIAGRAM³:



Compatibility DIAGRAM of $\hat{\circ}$ with $\hat{\subseteq}$

Furthermore, composition via pullback above then is associative, by *natural equivalence* of the (finite) limits defining compositions

$$h \hat{\circ} (g \hat{\circ} f), (h \hat{\circ} g) \hat{\circ} f : A \rightarrow B \rightarrow C \rightarrow D.$$

³F. Hermann


Cylindrification is componentwise, and gives the Cartesian Product for $\mathbf{PR}_A$ as a *monoidal* – again bifunctorial one – within extended Theory $\widehat{\mathbf{PR}}_A \sqsupset \mathbf{PR}_A$. But this extended Product does not have anymore (GODEMENT's) universal properties of a *Cartesian* Product, within $\widehat{\mathbf{PR}}_A$.

Iteration in $\widehat{\mathbf{PR}}_A$ works analogously to composition, using in this case *pullback iteration*.

Equality Definability: There is such a Theorem also for *partial map* Theory $\widehat{\mathbf{PR}}_A$ and its strengthenings.

Structure Theorem for $\widehat{\mathbf{S}}$:

- (i) $\widehat{\mathbf{S}}$ carries a – canonical – structure of a *diagonal symmetric monoidal category*, with *partial* composition $\widehat{\circ}$ and identities introduced above, (monoidal) product $\times$ extending $\times$ of $\mathbf{S}$, *association* $\text{ass} : (A \times B) \times C \xrightarrow{\cong} A \times (B \times C)$, *symmetry* (“*transposition*”) $\Theta : A \times B \xrightarrow{\cong} B \times A$, and *diagonal* $\Delta : A \rightarrow A \times A$ inherited from $\mathbf{S}$; cf. BUDACH & HOEHNKE 1975 and – later⁴ – PFENDER 1974 for an axiomatic approach to categories with a given type of *substitution* transformations. Our present theory $\widehat{\mathbf{S}}$, a theory of *partial* PR maps, is a monoidal category, which has – in addition to *natural* transformations ass , Θ , and Δ above –, so-called *half-terminal* maps, and the former projections as *half-projective* ones, in the terminology of BUDACH & HOEHNKE, “*half*” since the latter *natural* families of Theory $\mathbf{S}$, are no longer *natural transformations* for Theory $\widehat{\mathbf{S}}$. All of this *substitutive* structure is obviously preserved by the embedding $\mathbf{S} \sqsubset \widehat{\mathbf{S}}$.
- (ii) The defining diagram for an $\widehat{\mathbf{S}}$ -map – namely



Partial Map DIAGRAM

– constitutes in fact a *commuting* $\widehat{\mathbf{S}}$ diagram.

Conversely – with same notation as above – **define** the *minimised opposite* to d_f , beginning with formally *partial*, $\widehat{\mathbf{S}}$ map

$$d_f'^- = \langle (d_f, [\]_{\widehat{f}}) : D_f \rightarrow A \times D_f \rangle : A \rightarrow D_f,$$

opposite (graph) to given $\mathbf{S}$ map $d_f : D_f \rightarrow A$. This opposite is made *right-unique* by selecting D_f -minimal $\widehat{f}$ equivalence representant

$$[\]_{\widehat{f}} = [\hat{a}]_{\widehat{f}} =_{\text{def}} \min_{D_f} \{ \hat{a}' \leq \hat{a} \mid \widehat{f}(\hat{a}') \doteq_B \widehat{f}(\hat{a}) \} : D_f \rightarrow D_f :$$

⁴there is an earlier preprint of BUDACH & HOEHNKE

Minimal with respect to here “canonical”, CANTOR-ordering on **S** Object $D_f = \{D \mid \delta : D \rightarrow 2\}$. This order is *inherited* from D_f ’s “mother” fundamental Object, D , say. This object in turn is (well) ordered via canonical *counting*

$$\text{cantor}_D = \text{cantor}_D(n) : \mathbb{N} \xrightarrow{\cong} D,$$

(see general schema above of *PR dominated minimum*), and **get** the commuting $\widehat{\mathbf{S}}$ -DIAGRAM

$$\begin{array}{ccc} D_f & \xrightarrow{\hat{f}} & B \\ d_f^- \uparrow & \cong & \uparrow d_f^- \\ A & \xrightarrow{f} & B \end{array} \quad \text{put together:} \quad \begin{array}{ccc} D_f & \xrightarrow{\hat{f}} & B \\ d_f^- \uparrow & \cong & \uparrow d_f^- \\ A & \xrightarrow{f} & B \end{array}$$

Basic Partial Map DIAGRAM

- (iii) The first factor $f : A \rightarrow B$ in a $\widehat{\mathbf{PR}}_{\mathbf{A}}$ -composition $h = g \hat{\circ} f : A \rightarrow B \rightarrow C$, when giving an (embedded) $\mathbf{PR}_{\mathbf{A}}$ map $h : A \rightarrow C$, is itself an (embedded) $\mathbf{PR}_{\mathbf{A}}$ map: *first factor of a total map is total*.

So each *section* (“coretraction”) of $\widehat{\mathbf{PR}}_{\mathbf{A}}$ is a $\mathbf{PR}_{\mathbf{A}}$ map, in particular a $\widehat{\mathbf{PR}}_{\mathbf{A}}$ section of a $\mathbf{PR}_{\mathbf{A}}$ map is in $\mathbf{PR}_{\mathbf{A}}$.

- (iv) $\widehat{\mathbf{S}}$ clearly inherits from **S** FOURMAN’s *uniqueness equation*:

$$\text{For } h : C \rightarrow A \times B \text{ in } \widehat{\mathbf{S}} : h \hat{=} (h \hat{\circ} \ell, h \hat{\circ} r) : C \rightarrow A \times B,$$

where for $f : C \rightarrow A$, $g : C \rightarrow B$,

$$(f, g) =_{\text{def}} (f \times g) \hat{\circ} \Delta_C : C \rightarrow C \times C \rightarrow A \times B,$$

with *diagonal* $\Delta_C : C \rightarrow C \times C$ of **S**.

This equation guarantees *uniqueness* of the “induced” $(f, g) : C \rightarrow A \times B$, but (f, g) does not satisfy (both of) the Cartesian equations

$$\ell \hat{\circ} (f, g) \hat{=} f \quad \text{and} \quad r \hat{\circ} (f, g) \hat{=} g,$$

except f and g have *equal domains of definition*, i.e. if $i : D_f \rightarrow D_g$, $j : D_g \rightarrow D_f$ are available such that $d_g \circ i = d_f : D_f \rightarrow A$ as well as $d_f \circ j = d_g : D_g \rightarrow A$.

- (v) *Iteration* $f^{\S} : A \times \mathbb{N} \rightarrow A$ of $\widehat{\mathbf{S}}$ -endo is available in $\widehat{\mathbf{S}}$, satisfying the characteristic $\widehat{\mathbf{S}}$ -equations

$$\begin{aligned} f^{\S} \hat{\circ} (\text{id}_A, 0) &=_{\text{by def}} f^{\S} \hat{\circ} (A \times 0) \hat{\circ} \Delta_A \hat{=} \text{id}_A : A \rightarrow A, \quad \text{and} \\ f^{\S} \hat{\circ} (A \times s) &\hat{=} f \hat{\circ} f^{\S} : A \times \mathbb{N} \rightarrow A. \end{aligned}$$

(vi) Freyd's uniqueness of the *initialised iterated* holds in $\widehat{\mathbf{S}}$:

$$\begin{array}{c}
 f : A \rightarrow B, \ g : B \rightarrow B, \ h : A \times \mathbb{N} \rightarrow B \text{ in } \widehat{\mathbf{S}} \text{ such that} \\
 h \hat{\circ} (\text{id}_A, 0) \rightarrow f : A \rightarrow B \text{ and } \\
 h \hat{\circ} (A \times s) \hat{=} g \hat{\circ} h : A \times \mathbb{N} \rightarrow B \\
 \hline
 \text{(FR!)}_{\widehat{\mathbf{S}}} \quad h \hat{=} g^{\S} \hat{\circ} (f \times \mathbb{N}) : A \times \mathbb{N} \rightarrow B \times \mathbb{N} \rightarrow B.
 \end{array}$$

[The latter two statements are not so easy to prove: PR *construction* of *comparison* maps is needed, for comparing the respective enumerations of defined arguments in the postcedent, proceeding from the comparison maps given by the antecedents.]

(vii) For extension $\widehat{\mathbf{S}}$ of $\mathbf{S}$ again, we get – by the general FREYD's argument – the corresponding *full schema of primitive recursion*, namely

$$\begin{array}{c}
 g : A \rightarrow B \text{ in } \widehat{\mathbf{S}} \text{ (initialisation),} \\
 h : (A \times \mathbb{N}) \times B \rightarrow B \text{ (step map)} \\
 \hline
 \text{(pr)}_{\widehat{\mathbf{S}}} \quad f = \text{pr}[g, h] : A \times \mathbb{N} \rightarrow B \text{ is available in } \widehat{\mathbf{S}}, \\
 \text{characterised (up to equality } \hat{=} \text{) in } \widehat{\mathbf{S}} \text{ by} \\
 f \hat{\circ} (\text{id}_A, 0) \hat{=} g : A \rightarrow B \text{ and } \\
 f \hat{\circ} (A \times s) \hat{=} h \hat{\circ} (\text{id}_{A \times \mathbb{N}}, f) \\
 =_{\text{by def}} h \hat{\circ} ((A \times \mathbb{N}) \times f) \hat{\circ} \Delta_{A \times \mathbb{N}} : \\
 A \times \mathbb{N} \rightarrow (A \times \mathbb{N})^2 \rightarrow (A \times \mathbb{N}) \times B \rightarrow B.
 \end{array}$$

The **Proof** of this **Structure Theorem** is long, already since we have to show that many assertions; but mainly since assertion (vi) needs some auxiliary arguments.

Nevertheless, all of these assertions look plausible: they are “straightforward” extrapolations from the case of *finite* partial maps, by means we have at our disposition for the *potentially infinite*, primitive recursive case as *basic* theory.⁵

5 Partial-Map Extension as a Closure Operator


Closure: Theory $\widehat{\widehat{\mathbf{S}}}$ of *partial* maps over $\widehat{\mathbf{S}}$, of *partial* partial maps over $\mathbf{S}$, is (category) equivalent to Theory $\widehat{\mathbf{S}}$. Theory $\mathbf{S}$ is a strengthening of $\mathbf{PR}_A$ as always here.

Mutatis mutandis, construction of *Partiality Hull* $\widehat{\mathbf{S}} \sqsupset \mathbf{S}$ above of a *Cartesian* PR theory $\mathbf{S}$ can be applied again to *diagonal monoidal* Theory $\widehat{\mathbf{S}}$ “again”.

⁵full Proof is ready within detailed version, as a pdf file

Because of lack of Cartesianness this is more involved, and so is verification of the properties of this *Double Closure* $\widehat{\widehat{\mathbf{S}}}$. In particular it is more difficult to define composition: If you want to go into this detail, look at next DIAGRAM:

For **defining** composition of such $\widehat{\widehat{\mathbf{S}}}$ -morphisms, composition of say $f : A \rightarrow B$ and $g : B \rightarrow C$, consider the following $\mathbf{S}/\widehat{\widehat{\mathbf{S}}}$ -DIAGRAM which displays the $\widehat{\widehat{\mathbf{S}}}/\mathbf{S}$ data of f and g to be *composed* into an $\widehat{\widehat{\mathbf{S}}}$ -morphism $g \widehat{\widehat{\circ}} f : A \rightarrow B \rightarrow C$:



Composition DIAGRAM for $\widehat{\widehat{\mathbf{S}}}$


Composition $g \widehat{\widehat{\circ}} f : A \rightarrow C$ then is **defined** to have as *graph* $\gamma_{g \widehat{\widehat{\circ}} f}$ the map “induced” by the left and right *frame* morphisms of the DIAGRAM, namely:

$$\gamma_{g \widehat{\widehat{\circ}} f} =_{\text{def}} (\ell \widehat{\circ} \gamma f \widehat{\circ} d_{\gamma f} \circ \pi_\ell, r \widehat{\circ} \gamma g \widehat{\circ} d_{\gamma g} \circ \pi_r) : D_{g \widehat{\widehat{\circ}} f} \rightarrow A \times C.$$

The next assertion (really) to be **proved** is *idempotence* of our *Closure* operator, namely that each $\widehat{\widehat{\mathbf{S}}}$ map $f : A \rightarrow B$ is represented – with respect to notion of equality $\widehat{\widehat{=}}$ of $\widehat{\widehat{\mathbf{S}}}$, by a suitable $\widehat{\widehat{\mathbf{S}}}$ -map $h : A \rightarrow B$.

For a **Proof** look at the following DIAGRAM, for given $\widehat{\widehat{\mathbf{S}}}$ -map

$$f = \langle \gamma f = (\ell \widehat{\circ} \gamma f, r \widehat{\circ} \gamma f) =: (d_f, \widehat{f}) : D_f \rightarrow A \times B \rangle : A \rightarrow B :$$



Closure DIAGRAM for Extension by partial maps

In this DIAGRAM, $\gamma f : D_f \rightarrow A \times B$ is the *graph* of $\widehat{\widehat{\mathbf{S}}}$ -morphism $f : A \rightarrow B$ to be considered. The $\mathbf{S}$ -maps $d_{\gamma f} : D_{\gamma f} \rightarrow D_f$ (defined-arguments enumeration) and $\widehat{\gamma f} : D_{\gamma f} \rightarrow A \times B$ (rule) are to define $\gamma f : D_f \rightarrow A \times B$ as a *partial* $\mathbf{S}$ -map, an $\widehat{\mathbf{S}}$ morphism.

This DIAGRAM shows the way of **Proof** for

Closure Theorem for Extension of Theory $\mathbf{S}$ by Partial Maps:

Closure by Partial Maps is idempotent: Partial map Closure of theory $\widehat{\mathbf{S}}$ is again a diagonal monoidal category $\widehat{\widehat{\mathbf{S}}}$ which is in fact *equivalent* – as such a category – to theory $\widehat{\mathbf{S}}$:

$$\widehat{\widehat{\mathbf{S}}} \cong \widehat{\mathbf{S}}.$$

[Both inherit – from $\mathbf{S}$ – Object $\mathbb{N}$ as NNO in the sense of the (full) schema of PR for $\mathbb{N}$.]

6 μ -Recursion without Quantifiers

Church type “Inclusion”: For given $\mathbf{PR}_A$ -predicate $\varphi : A \times \mathbb{N} \rightarrow 2$, *partially defined* “map”

$$\mu\varphi = \mu\varphi(a) : A \rightarrow \mathbb{N},$$

classically defined by

$$\mu\varphi(a) = \begin{cases} \min\{n \in \mathbb{N} \mid \varphi(a, n)\} & \text{if } (\exists n \in \mathbb{N}) \varphi(a, n) \\ \text{undefined} & \text{if } (\forall n \in \mathbb{N}) \neg \varphi(a, n), \end{cases}$$

has a – classically *correct – representation* within (strengthenings of) Theory $\widehat{\mathbf{PR}}_{\mathbf{A}}$ as

$$\begin{array}{ccc}
 D_{\mu\varphi} =_{\text{def}} \{A \times \mathbb{N} \mid \varphi\} = \{(a, n) \mid \varphi(a, n)\} & & \\
 \downarrow d_{\mu\varphi} = \ell \circ \subseteq & \searrow \widehat{\mu\varphi} & \\
 A & \xrightarrow{\mu\varphi} & \mathbb{N},
 \end{array}$$

Here *defined-arguments* (PR) enumeration is

$$d_{\mu\varphi} = d_{\mu\varphi}(a, n) =_{\text{def}} a : \{A \times \mathbb{N} \mid \varphi\} \subseteq A \times \mathbb{N} \xrightarrow{\ell} A,$$

and rule $\widehat{\mu\varphi} : \{(a, n) \mid \varphi(a, n)\} \rightarrow \mathbb{N}$ is (totally) PR defined by

$$\widehat{\mu\varphi} = \widehat{\mu\varphi}(a, n) = \min\{m \leq n \mid \varphi(a, m)\} : \{(a, n) \mid \varphi(a, n)\} \rightarrow \mathbb{N}.$$


μ -Lemma: $\widehat{\mathbf{S}}$ admits the following (Free-Variables) schema (μ) , operator μ 's “property”, combined with *uniqueness* schema $(\mu!)$, as a *characterisation* of the μ -operator $\langle \varphi : A \times \mathbb{N} \rightarrow 2 \rangle \mapsto \langle \mu\varphi : A \rightarrow \mathbb{N} \rangle$ above:

$$\begin{array}{l}
 \varphi = \varphi(a, n) : A \times \mathbb{N} \rightarrow 2 \text{ } \mathbf{S}\text{-map (‘‘predicate’’),} \\
 [a \in A \text{ (free) is the } \textit{passive} \text{ parameter,} \\
 n \in \mathbb{N} \text{ free the } \textit{recursion parameter}] \\
 (\mu) \quad \hline
 \mu\varphi = \langle (d_{\mu\varphi}, \widehat{\mu\varphi}) : D_{\mu\varphi} \rightarrow A \times \mathbb{N} \rangle : A \rightarrow \mathbb{N} \\
 \text{is an } \widehat{\mathbf{S}}\text{-map such that} \\
 \mathbf{S} \vdash \varphi(d_{\mu\varphi}(\hat{a}), \widehat{\mu\varphi}(\hat{a})) = \text{true}_{D_{\mu\varphi}} : D_{\mu\varphi} \rightarrow 2, \\
 [\hat{a} \in D_{\mu\varphi} \text{ free, so just } a \in A \text{ of form } a := d_{\mu\varphi}(\hat{a}) \\
 \text{counts as – is } \textit{enumerated} \text{ as – ‘‘defined argument’’ for } \mu\varphi] \\
 + \text{ ‘‘argumentwise’’ } \mathbf{minimality}: \\
 \mathbf{S} \vdash [\varphi(d_{\mu\varphi}(\hat{a}), n) \implies \widehat{\mu\varphi}(\hat{a}) \leq n] : D_{\mu\varphi} \times \mathbb{N} \rightarrow 2
 \end{array}$$

as well as *uniqueness*, by *maximal extension*:

$$\begin{array}{l}
 f = f(a) : A \rightarrow \mathbb{N} \text{ in } \widehat{\mathbf{S}} \text{ such that} \\
 \mathbf{S} \vdash \varphi(d_f(\hat{a}), \widehat{f}(\hat{a})) = \text{true}_{D_f} : D_f \rightarrow 2, \\
 (\mu!) \quad \mathbf{S} \vdash \varphi(d_f(\hat{a}), n) \implies \widehat{f}(\hat{a}) \leq n : D_f \times \mathbb{N} \rightarrow 2 \\
 \hline
 \mathbf{S} \vdash f \subseteq \mu\varphi : A \rightarrow \mathbb{N} \text{ (inclusion of graphs).}
 \end{array}$$

Proof of schema (μ) is by **definition** of $\mu\varphi = \mu\varphi(a) : A \rightarrow \mathbb{N}$. **Proof idea** for uniqueness schema $(\mu!)$ is “displayed” as the following DIAGRAM:



μ -applied-to-**S**-predicates DIAGRAM gives $f \hat{\subseteq} \mu\varphi$.

Remark: Within PEANO-Arithmétique **PA**, and hence also within set theory, our $\mu\varphi : A \rightarrow \mathbb{N}$ equals

$$\mu\varphi = \langle (\subseteq, \hat{\mu}\varphi) : \hat{A} \rightarrow A \times \mathbb{N} \rangle : A \supset \hat{A} \rightarrow \mathbb{N},$$

with $\hat{A} = \{\hat{a} \in A \mid \exists n \varphi(\hat{a}, n)\}$, and $\hat{\mu}\varphi(\hat{a}) = \min\{m \in \mathbb{N} \mid \varphi(\hat{a}, m)\} : \hat{A} \rightarrow \mathbb{N}$, i.e. it is given there by the classical *partial* minimum definition. But this definition lacks *constructivity*, since $\hat{A} \subseteq A$ is in general not PR decidable.

Conversely, consider a *partial* PR map,

$$f = \langle (d_f, \hat{f}) : D_f \rightarrow A \times B \rangle : A \rightarrow B, \text{ out of } \mathbf{PR}_{\mathbf{A}}.$$

Standard, pointed case: f is *defined* at least at one *point*, say at $a_0 = d_f(\hat{a}_0) : \mathbb{1} \rightarrow D_f \rightarrow A$.

Such f is represented easily, within Theory $\mathbf{PR}_{\mathbf{A}}$, by a μ -recursive $\mathbf{PR}_{\mathbf{A}}$ -map (followed by a $\mathbf{PR}_{\mathbf{A}}$ map), namely by

$$\mu[f] \stackrel{\text{def}}{=} (\hat{f} \circ \text{count}_{D_f}) \hat{\circ} \mu\varphi_f : A \rightarrow \mathbb{N} \rightarrow D_f \rightarrow \mathbb{N}.$$

$\mathbf{PR}_{\mathbf{A}}$ -predicate $\varphi_f = \varphi_f(a, n) : A \times \mathbb{N} \rightarrow 2$ is given as

$$\varphi_f = \varphi_f(a, n) \stackrel{\text{def}}{=} [a \dot{=}_A d_f \circ \text{count}_{D_f}(n)] : A \times \mathbb{N} \rightarrow A \times D_f \rightarrow 2 \text{ PR}.$$

Here $\text{count}_{D_f} : \mathbb{N} \rightarrow D_f$ designates a (retractive) (PR) *count* of D_f . For disposing on this *count* of D_f , we had to assume that D_f comes with a $\mathbf{PR}_{\mathbf{A}}$ -*point*, $\hat{a}_0 : \mathbb{1} \rightarrow D_f$ above.

Partial map $\mu\varphi_f : A \rightarrow \mathbb{N}$ is the genuine μ -recursive kernel of μ -representation $\mu[f] : A \rightarrow B$ of (pointed) partial map $f : A \rightarrow B$. We count *composition* of μ -recursive maps with PR maps equally under the μ -recursive ones. So in this sense, $\mu[f] \hat{=} f : A \rightarrow B$ is a μ -recursive representant of f within $\mathbf{PR}_{\mathbf{A}}$ and its strengthenings, a (partial) map in $\mu\mathbf{R}$, and in $\mu\mathbf{S}$ for strengthenings **S** of $\mathbf{PR}_{\mathbf{A}}$.

The case that D_f has *no point*, and is nevertheless *not S-derivably* empty, causes a **problem**, formally. We “solve” this problem by modifying (extending) the definition of $\mu[f] : A \rightarrow \mathbb{N}$ as follows:

PR_A-Object D_f is predicative restriction $D_f : D \rightarrow 2$ of a *fundamental* Object D , which comes as such with a (componentwise) zero $0 : \mathbb{1} \rightarrow D$ as (privileged) *point*. This Object admits a CANTOR count $\text{count}_D : \mathbb{N} \xrightarrow{\cong} D$. (Trivial exception: for $D = \mathbb{1}$, $\text{count}_D = ! : \mathbb{N} \rightarrow \mathbb{1}$ is still a retraction.)

In present general case, replace – on the way – Object $A \subset \mathbb{X}$ by sum, (disjoint) union $\mathbb{1} \oplus A \subset \mathbb{X}$, and define $\dot{\varphi}_f = \dot{\varphi}_f(a, n) : A \times \mathbb{N} \rightarrow 2$ by the following **PR_A**-DIAGRAM:

$$\begin{array}{ccc}
A \times \mathbb{N} & \xrightarrow{\dot{\varphi}_f} & 2 \\
\downarrow \iota \times \mathbb{N} \subset & & \uparrow \doteq \\
(\mathbb{1} \oplus A) \times \mathbb{N} & \stackrel{=_{\text{def}}}{=} & (\mathbb{1} \oplus A) \times (\mathbb{1} \oplus A) \\
\downarrow (\mathbb{1} \oplus A) \times \text{count}_D & & \uparrow \text{id} \times (!_{D \setminus D_f} \oplus d_f) \\
(\mathbb{1} \oplus A) \times D & \stackrel{=}{=} & (\mathbb{1} \oplus A) \times ((D \setminus D_f) \oplus D_f)
\end{array}$$

$\dot{\varphi}_f = \dot{\varphi}_f(a, n) : A \times \mathbb{N} \rightarrow 2$ in Free-Variables notation:

$$\begin{aligned}
A \times \mathbb{N} \ni (a, n) &\stackrel{\iota}{\mapsto} (a, n) \mapsto (a, \text{count}_D(n)) \\
&\mapsto \begin{cases} \text{false} & \text{if } \text{count}_D(n) \notin D_f, \text{ (“outside” case),} \\ [a \doteq_A d_f(\text{count}_D(n))] [\in 2] & \text{if } \text{count}_D(n) \in D_f. \end{cases}
\end{aligned}$$

This given, we **define** in this general case

$$\mu[f] \stackrel{=_{\text{def}}}{=} \widehat{f} \circ \text{count}_D \circ \mu \dot{\varphi}_f : A \rightarrow \mathbb{N} \rightarrow D \rightarrow B.$$

Note first that $\widehat{f} : D_f \rightarrow B$ comes by (formal) Domain restriction of a genuine **PR** map $\widehat{f} : D \rightarrow B' : D_f = \{D \mid D_f\}$, D (and B') fundamental: This by definition of *maps* of Theory **PR_A** = **PR** + (abstr).

Second: wider count count_D , available in particular for D as a fundamental Object, Codomain-restricts here nicely, gives “again” μ -representation of f , here via $D \supseteq D_f = \{D \mid D_f : D \rightarrow 2\}$.

This taken together gives μ -representation of general partial PR-map $f : A \rightarrow B$ as

$$f \hat{=} \mu[f] \stackrel{=_{\text{by def}}}{=} (\widehat{f} \circ \text{count}_D) \circ \mu \dot{\varphi}_f : A \xrightarrow{\mu \dot{\varphi}_f} \mathbb{N} \xrightarrow{\text{count}_D} D \xrightarrow{\widehat{f}} B.$$

So we have reached

Another Proved Instance of Church’s Thesis:

- The notion of a μ -recursive (partial) map is *equivalent* to that of a *Partial* PR map, over “all” Theories of Primitive Recursion.

- Theories $\widehat{\mathbf{PR}}_{\mathbf{A}}$ and $\mu\mathbf{R}$ are equivalent, and the **Closure Theorem** $\widehat{\widehat{\mathbf{S}}} \cong \widehat{\mathbf{S}}$ above then shows $\mu\mu\mathbf{R} \cong \mu\mathbf{R}$:
- *Level-one μ -recursion is enough* for getting all μ -recursive maps. By the above, this gives the well known corresponding result for while programs: *one while loop is enough*: Any such program can be equivalently transformed into a while loop program without nesting of while loops.
- All this works as well for *strengthenings* $\mathbf{S}$, $\widehat{\mathbf{S}}$ of $\mathbf{PR}_{\mathbf{A}}$ and $\widehat{\mathbf{PR}}_{\mathbf{A}}$ respectively. We would name the corresponding Theory of μ -recursion $\mu\mathbf{S} \cong \widehat{\mathbf{S}}$.

Conclusion so far:

- We can *eliminate formal existential quantification* – as well as (individual, formal) *variables* – from the theory of μ -recursion: we interpret application of μ -operator to predicates of theories $\mathbf{S}$ strengthening PR Theory $\mathbf{PR}_{\mathbf{A}} = \mathbf{PR} + (\text{abstr})$ as suitable *partial* maps, maps in Theory $\widehat{\mathbf{S}}$.
- The μ -operator canonically extends to all *partial* predicates $\varphi : A \times \mathbb{N} \rightarrow 2$, and associates to them just partial maps $\mu\varphi : A \rightarrow \mathbb{N}$, within $\widehat{\mathbf{S}}$ itself. So, “once again”, we see, that theories $\widehat{\mathbf{S}}$ of *partial PR maps* are *closed*, this time under the μ -operator, “in parallel” to *Closure* of $\widehat{\mathbf{S}}$ under forming *partial* maps: *partial partial PR maps* “are” *partial PR maps*.
- We have the following chain of categorical *equivalences* of theories considered so far:

$$\mathbf{S} \sqsubset \mu\mathbf{S} \cong \mu\mu\mathbf{S} \cong \widehat{\widehat{\mathbf{S}}} \cong \widehat{\mathbf{S}} \sqsupset \mathbf{S},$$

the inclusions being *diagonal-monoidal PR compatible* with the equivalences.

[A *partial* PR map $f : A \rightarrow B$ which is, “by hazard”, a *total* map – discussion of overall *termination = total definedness* in part RCF 2 ($\varepsilon\&\mathcal{C}$), is in general *not* itself PR: only its graph $(d_f, f) : D_f \rightarrow A \times B$ is PR. ACKERMANN type maps, in particular *evaluation* of all PR-map-codes, are formally partial maps. In *well defined cases*, they can be *forced* – by plausible additional **axiom** – to become *on-terminating*, i.e. to get defined-argument enumeration epimorphic.]

- Conversely, application of the μ -operator, already just to $\mathbf{PR}_{\mathbf{A}}$ -predicates $\varphi = \varphi(a, n) : A \times \mathbb{N} \rightarrow 2$, *generates* all *partial* $\mathbf{PR}_{\mathbf{A}}$ -maps $f : A \rightarrow B$.
- As important special cases of basic PR theories $\mathbf{S}$ we have at the moment Theory $\mathbf{PR}_{\mathbf{A}} = \mathbf{PR} + (\text{abstr})$ itself as well as the PR *trace* $\mathbf{PA} \upharpoonright \mathbf{PR}_{\mathbf{A}}$ of $\mathbf{PA}$: All $\mathbf{PR}_{\mathbf{A}}$ -maps (map terms) with all those equations in between, which are derivable by $\mathbf{PA}$: Our theories, notions, and results have a structure-preserving Interpretation into (within) Peano-Arithmetic $\mathbf{PA}$.

Same for **set theories** in place of **PA**, in particular **ZF**, **ZFC** and their first order subsystems **1ZF**, **1ZFC** = **1ZF** + **ACC**. **ACC** is the Axiom of *Countable Choice*.

7 Content Driven Loops, in particular while Loops

Classically, *with* variables, a while loop $\text{wh} = \text{wh}[\chi | f]$ is “defined” in *pseudocode* by

$$\begin{aligned} \text{wh}(a) &:= [a' := a; \\ &\quad \text{while } \chi(a') \text{ do } a' := f(a') \text{ od}; \\ &\quad \text{wh}(a) := a']. \end{aligned}$$

The formal version of this – within a *classical*, element based setting – is the following partial-(PEANO)-map characterisation:

$$\text{wh}(a) = \text{wh}[\chi | f](a) = \begin{cases} a & \text{if } \neg \chi(a) \\ \text{wh}(f(a)) & \text{if } \chi(a) \end{cases} : A \rightarrow A.$$

It is possible to give a *static Definition* of $\text{wh} = \text{wh}[\chi | f] : A \rightarrow A$, within $\widehat{\mathbf{PR}}_{\mathbf{A}} \sqsubset \mathbf{PR}_{\mathbf{A}}$ (and strengthenings) as follows:

$$\begin{aligned} \text{With } \varphi &= \varphi_{[\chi | f]}(a, n) \stackrel{\text{def}}{=} \neg \chi f^n(a) \\ &\stackrel{\text{by def}}{=} \neg \chi f^{\S}(a, n) : A \times \mathbb{N} \rightarrow A \rightarrow 2 \rightarrow 2, \\ \text{the } \underline{\text{while}} \text{ loop} \\ \text{wh} &= \text{wh}[\chi | f] : A \rightarrow A \text{ is given as} \\ \text{wh} &\stackrel{\text{def}}{=} f^{\mu \varphi_{[\chi | f]}}(a) \\ &\stackrel{\text{by def}}{=} f^{\S} \hat{\circ} (\text{id}_A, \mu \varphi_{[\chi | f]}) \\ &\stackrel{\text{by def}}{=} f^{\S} \hat{\circ} (A \times \mu \varphi_{[\chi | f]}) \hat{\circ} \Delta_A : \\ &A \rightarrow A \times A \rightarrow A \times \mathbb{N} \rightarrow A. \end{aligned}$$

In this generalised categorical sense, we have within theories $\widehat{\mathbf{S}} \sqsubset \mathbf{S}$ ($\mathbf{S}$ strengthening $\mathbf{PR}_{\mathbf{A}}$), all while loops.

Characterisation Theorem for while loops *over* $\mathbf{S}$, within Theory $\widehat{\mathbf{S}}$: For $\chi : A \rightarrow 2$ (*control*) and $f : A \rightarrow A$ (*step*), both – for the time being – $\mathbf{S}$ -maps, while loop $\text{wh} = \text{wh}[\chi | f] : A \rightarrow A$ (as defined above), is characterised by the following *implications* within $\widehat{\mathbf{S}}$:

$$\begin{aligned} \widehat{\mathbf{S}} &\vdash \neg \chi \circ a \implies \text{wh} \hat{\circ} a \doteq a : A \rightarrow 2, \text{ uland} \\ \widehat{\mathbf{S}} &\vdash \chi \circ a \implies \text{wh} \hat{\circ} a \doteq \text{wh} \hat{\circ} f \circ a. \end{aligned}$$

Dominated Termination of while loop

$$\text{wh} = \text{wh}[\chi \mid f] = \text{wh}[\chi \mid f](a) : A \rightarrow A,$$

at argument $a \in A$, is expressed by

$$[m \text{ def wh}[\chi \mid f](a)] =_{\text{def}} \neg \chi f^m(a) : \mathbb{N} \times A \rightarrow 2 :$$

In words: *iteration of endo $f : A \rightarrow A$, applied to argument a , reaches stop condition $\neg \chi$ after (at most) m steps.* Here both: argument $a \in A$ as well as iteration counter $m \in \mathbb{N}$, are free variables (categorically: projections). $[m \text{ def wh}[\chi \mid f](a) \doteq_A \bar{a}]$ is to mean:

$$\begin{aligned} & [m \text{ def wh}[\chi \mid f](a)] \\ & \wedge \text{wh}[\chi \mid f](a) \doteq_A f^{\min\{n \leq m \mid \neg \chi f^n(a)\}}(a) \\ & =_{\text{by def}} f^{\S}(a, \min\{n \leq m \mid \neg \chi f^n(a)\}) \doteq_A \bar{a} : \\ & \mathbb{N} \times A^2 \rightarrow 2, \ m \in \mathbb{N}, \ a, \bar{a} \in A \text{ free.} \end{aligned}$$

[“Variable” $n \in \mathbb{N}$ used in the min PR Operator is *auxiliary, bounded* by m . It does not count as a (free) variable.]

From a *logical* point of view, there are – at least – the following open **Questions**, in

Arithmetics Complexity Problem:

- (i) Does Theory **PR** admit *strict, consistent* strengthenings, or is it a *simple theory*, will say that it admits its given notion of equality and the indiscrete (inconsistency) equality of its maps as only “congruences?”, cf. a simple *group* which has as *normal subgroups* only itself and $\{1\}$. Because of reasons to be explained in later work, my guess here is: **PR** *admits* non-trivial strengthenings, in particular I suppose that the PR *trace* of **PA**, explained above, is a strict strengthening of **PR** resp. **PRA** = **PR** + (abstr).

We cannot exclude at present that all these strengthenings of **PR** make up a whole lattice of (Free-Variables) Arithmetical Theories, each of them giving particular, “new” features to Primitive Recursive Arithmetics.

- (ii) Already at start we possibly have such a strengthening: If Free-Variables (“Free Variables” in the classical sense) *Primitive Recursive Arithmetic* **PRA** is **defined** to have as its terms all map terms obtainable by the (full) schema of Primitive Recursion, and as formulae just the *defining equations* for the maps introduced by that schema, then I see no way to **prove** all of the usual semiring equations for $\mathbb{N}$:

We *need* Freyd’s **uniqueness** (FR!) – section 1 above – of the *initialised iterated*: From this HORN clause we can show (!) in particular GOODSTEIN’s uniqueness rules U_1 to U_4 on which *his* Proof of the semiring properties of $\mathbb{N}$ is based. He takes these rules as **axioms**.

My guess is here – if I have understood right the definition of **PRA** – that **PR** = **PRA** + (FR!) is a strict strengthening of **PRA**, at least if there is no “underground” connection to the set theoretic view of maps as (possibly infinite) *argument-value tables*.

- (iii) So again, Arithmetic would *become* simpler, if Theory **PR** would turn out to be *simple*. If not, we have a diversity of Arithmetics, a diversity intuitively far below such issues as Independence of the Axiom of Choice or of the Continuum Hypothesis. At least the latter is open in the context of a **Constructive Analysis** based on map theoretic, Free-Variables (variable-free) Primitive Recursive and μ -recursive Arithmetics.

References

- J. BARWISE ed. 1977: *Handbook of Mathematical Logic*. North Holland.
- H.-B. BRINKMANN, D. PUPPE 1969: *Abelsche und exakte Kategorien, Korrespondenzen*. L.N. in Math. **96**. Springer.
- L. BUDACH, H.-J. HOEHNKE 1975: *Automaten und Funktoren*. Akademie-Verlag Berlin.
- C. EHRESMANN 1965: *Catégories et Structures*. Dunod Paris.
- H. EHRLIG, W. KÜHNEL, M. PFENDER 1975: Diagram Characterization of Recursion. LN in Comp. Sc. 25, 137-143.
- S. EILENBERG, C. C. ELGOT 1970: *Recursiveness*. Academic Press.
- S. EILENBERG, S. MAC LANE 1945: General Theory of Natural Equivalences. *Trans. AMS* 58, 231-294.
- P. J. FREYD 1972: Aspects of Topoi. *Bull. Australian Math. Soc.* **7**, 1-76.
- R. L. GOODSTEIN 1971: *Development of Mathematical Logic*, ch. 7: Free-Variable Arithmetics. Logos Press.
- F. HAUSDORFF 1908: Grundzüge einer Theorie der geordneten Mengen. *Math. Ann.* **65**, 435-505.
- D. HILBERT: Mathematische Probleme. Vortrag Paris 1900. *Gesammelte Abhandlungen*. Springer 1970.
- P. T. JOHNSTONE 1977: *Topos Theory*. Academic Press
- A. JOYAL 1973: Arithmetical Universes. Talk at Oberwolfach.
- J. LAMBEK, P. J. SCOTT 1986: *Introduction to higher order categorical logic*. Cambridge University Press.
- F. W. LAWVERE 1964: An Elementary Theory of the Category of Sets. *Proc. Nat. Acad. Sc. USA* **51**, 1506-1510.
- F. W. LAWVERE, S. H. SCHANUEL 1997: *Conceptual Mathematics, A first introduction to categories*. Cambridge University Press.
- S. MAC LANE 1972: *Categories for the working mathematician*. Springer.
- B. PAREIGIS 1969: *Kategorien und Funktoren*. Teubner.
- R. PÉTER 1967: *Recursive Functions*. Academic Press.
- M. PFENDER 1974: Universal Algebra in S-Monoidal Categories. Algebra-Berichte Nr. 20, Mathematisches Institut der Universität München. Verlag Uni-Druck München.
- M. PFENDER 2008 RCF1d: Theories of PR Maps and Partial PR Maps, detailed version. pdf file.
- M. PFENDER, M. KRÖPLIN, D. PAPE 1994: Primitive Recursion, Equality, and a Universal Set. *Math. Struct. in Comp. Sc.* **4**, 295-313.
- H. REICHEL 1987: *Initial Computability, Algebraic Specifications, and Partial Algebras*. Oxford Science Publications.
- W. RAUTENBERG 1995/2006: *A Concise Introduction to Mathematical Logic*. Universitext Springer 2006.

R. REITER 1980: Mengentheoretische Konstruktionen in arithmetischen Universen. Diploma Thesis. Techn. Univ. Berlin.

R. REITER 1982: Ein algebraisch-konstruktiver Abbildungskalkül zur Fundierung der elementaren Arithmetik. Dissertation, rejected by the Mathematics Department of TU Berlin.

L. ROMÀN 1989: Cartesian categories with natural numbers object. *J. Pure and Appl. Alg.* **58**, 267-278.

W. W. TAIT 1996: Frege versus Cantor and Dedekind: on the concept of number. Frege, Russell, Wittgenstein: *Essays in Early Analytic Philosophy (in honor of Leonhard Linsky)* (ed. W. W. Tait). Lasalle: Open Court Press (1996): 213-248. Reprinted in *Frege: Importance and Legacy* (ed. M. Schirn). Berlin: Walter de Gruyter (1996): 70-113.

U. THIEL 1982: Der Körper der rationalen Zahlen im arithmetischen Universum. Diploma Thesis. Techn. Univ. Berlin.

A. TARSKI, S. GIVANT 1987: *A formalization of set theory without variables*. AMS Coll. Publ. vol. 41.

Address of the author:

M. PFENDER

Institut für Mathematik

Technische Universität Berlin

D-10623 Berlin

pfender@math.TU-Berlin.DE